arXiv:1303.4038v1 [hep-th] 17 Mar 2013

Striking Properties of Finite Nonrenormalizable Theories

Kevin Cahill*

Department of Physics & Astronomy,
University of New Mexico, Albuquerque, NM 87131, USA and
Physics Department, Fudan University, Shanghai 200433, China

(Dated: March 19, 2013)

Abstract

Some nonrenormalizable theories have remarkable properties. Their euclidian and minkowskian Green's functions are finite, and the features that make them finite may help us understand dark energy, confinement, and quantum gravity.

PACS numbers: 11.10.-z, 11.10.Lm, 11.15.Ha, 11.15.Tk, 12.38.Aw, 04.60.-m

---

* cahill@unm.edu

The existence of quantum theories with finite Green's functions was first suggested by Boettcher and Bender [1]. A recent paper [2] shows that a theory of a scalar boson with euclidian action density

$$L_e = \frac{1}{2}\left(\partial_\mu \phi(x)\right)^2 + \frac{1}{2}\, m^2 M^2 \, \frac{1}{1 - \phi^2(x)/M^2} \tag{1}$$

has euclidian Green's functions

$$G_e(x_1, \ldots, x_n) \equiv \langle 0|\mathcal{T}\left[\phi_e(x_1)\ldots\phi_e(x_n)\right]|0\rangle = \frac{\displaystyle\int \phi(x_1)\ldots\phi(x_n)\, \exp\left[-\int L_e(\phi)\, d^4x\right] D\phi}{\displaystyle\int \exp\left[-\int L_e(\phi)\, d^4x\right] D\phi} \tag{2}$$

that are finite. The Green's functions of the corresponding theory in minkowski space with action density

$$L_m = -\frac{1}{2}\,\partial_\mu\phi(x)\,\partial^\mu\phi(x) - \frac{1}{2}\, m^2 M^2 \, \frac{1}{1 - \phi^2(x)/M^2} \tag{3}$$

also are finite [2]. The somewhat more interesting theory with euclidian action density

$$L_{se} = \frac{1}{2}\left(\partial_\mu \phi(x)\right)^2 + m^2 M^2 \, \frac{1}{\sqrt{1 - \phi^2(x)/M^2}} \tag{4}$$

also has finite Green's functions (and also describes particles of mass m). These theories are not renormalizable, but one may use lattice methods and expansions in powers of $\hbar$ to extract physical information from them.

The derivatives in L_{se} dampen wiggles in the field $\phi(x)$ but may not drastically affect its mean value. If we ignore them, then the mean value of the potential-energy density in the ground state of the theory is a ratio of products of integrals all but one of which cancel

$$\begin{aligned}\langle 0|\frac{m^2M^2}{\sqrt{1-\phi^2(x)/M^2}}|0\rangle &= \frac{\displaystyle\int \frac{m^2M^2}{\sqrt{1-\phi^2(x)/M^2}}\,\exp\left\{-\int \frac{m^2M^2}{\sqrt{1-\phi^2(x')/M^2}}\,d^4x'\right\} D\phi}{\displaystyle\int \exp\left\{-\int \frac{m^2M^2}{\sqrt{1-\phi^2(x')/M^2}}\,d^4x'\right\} D\phi} \\ &= \frac{\displaystyle\int \frac{m^2M^2}{\sqrt{1-\phi^2(x)/M^2}}\,\exp\left\{-\frac{m^2M^2}{\sqrt{1-\phi^2(x)/M^2}}\,d^4x\right\} d\phi(x)}{\displaystyle\int \exp\left\{-\frac{m^2M^2}{\sqrt{1-\phi^2(x)/M^2}}\,d^4x\right\} d\phi(x)}.\end{aligned} \tag{5}$$

Setting $d^4x = \epsilon^4$ and taking the limit $\epsilon \to 0$, we find

$$m^2M^2\,\langle 0|\frac{1}{\sqrt{1-\phi^2(x)/M^2}}|0\rangle = 1.5708\; m^2M^2. \tag{6}$$

In the same approximation, the mean value of the potential-energy density of the first theory (1) diverges logarithmically.

The full energy density, of course, involves the derivatives. We can make the mean value of their contribution finite by using as the euclidian action density

$$L_{re} = M^4 \left\{1 - M^{-4} \left[(\partial_\mu \phi(x))^2 + m^2 \phi^2(x)\right]\right\}^{-1/2} . \tag{7}$$

We could presumably adjust the masses m and M so as to fit the experimental value of the dark-energy density [3]

$$\langle 0|L_{re}|0\rangle = \frac{\displaystyle\int L_{re} \, \exp\left[-\int L_{re}(\phi)\, d^4x\right] D\phi}{\displaystyle\int \exp\left[-\int L_{re}(\phi)\, d^4x\right] D\phi} = 3.1 \times 10^{-47} \text{ GeV}^4. \tag{8}$$

Many nonrenormalizable theories have Green's functions that are finite or less singular than those of a free field. We can make a typical theory of scalar and vector bosons less singular by replacing its euclidian action density L_e by [4]

$$L_e' = \frac{L_e}{1 - L_e/m^4} \tag{9}$$

or by

$$L_e'' = \frac{m^4}{\sqrt{1 - L_e/m^4}} \tag{10}$$

in which m is a suitable mass.

The euclidian action density of $SU(3)$ gauge theory is the trace

$$L_3 = \frac{1}{2g^2} \operatorname{Tr}\left(F_{\mu\nu}^2\right) \tag{11}$$

in which the Faraday matrix is $F_{\mu\nu} = g\, t^a F^a_{\mu\nu}$, the generators t^a of $SU(3)$ are half the Gell-Mann matrices, and $F^a_{\mu\nu} = \partial_\mu A^a_\nu - \partial_\nu A^a_\mu - g f_{abc} A^b_\mu A^c_\nu$. The theory described by

$$L_3' = \frac{m^4}{\sqrt{1 - L_3/m^4}} \tag{12}$$

has Green's functions that are less singular. And if the mean values in the vacuum of euclidian-time-ordered products of gauge-invariant operators O_k are given by the path integrals

$$\langle 0|\mathcal{T}\left[O_{1e}(x_1)\ldots O_{ne}(x_n)\right]|0\rangle = \frac{\displaystyle\int O_1(x_1)\ldots O_n(x_n)\, \exp\left[-\int L_3'\, d^4x\right] D\phi}{\displaystyle\int \exp\left[-\int L_3'\, d^4x\right] D\phi}, \tag{13}$$

then they are less singular than those of L_3 and may be finite.

Wilson [5], Creutz [6], and others have demonstrated quark confinement on the lattice by replacing the euclidian action of pure continuum QCD

$$S_{QCD} = \frac{1}{2g^2} \int \mathrm{Tr}\left(F_{\mu\nu}^2\right) \, d^4x = \int L_3 \, d^4x \tag{14}$$

by a sum over the plaquettes of a lattice

$$S_W = \sum_{\square} S_{\square} \tag{15}$$

of Wilson's action $S_{\square}$ which is the trace of the product of the elements U of $SU(3)$ on the links that surround the plaquette

$$S_{\square} = \beta \, \left[1 - (1/3)\mathrm{Re}\,\mathrm{Tr}\left(U_{ij}U_{jk}U_{k\ell}U_{\ell i}\right)\right]. \tag{16}$$

Yet there is a big difference between the continuum action density $\mathrm{Tr}(F_{\mu\nu}^2)/2g^2$ which can be arbitrarily large and Wilson's action $S_{\square}$ which is bounded by β. This gap is bridged if one uses the action density L_3' which keeps $\mathrm{Tr}(F_{\mu\nu}^2)/2g^2$ bounded like Wilson's $S_{\square}$.

The euclidian action density of general relativity is not bounded below, and so the recipes (9) and (10) don't work for it. Instead, we can use, for instance,

$$L_E' = \frac{1}{16\pi G_N^2} \left(\frac{\cosh^2\theta}{1 - G_N \, R_E} + \frac{\sinh^2\theta}{1 + G_N \, R_E} \right) \sqrt{|g|} \tag{17}$$

in which R_E is the euclidian Ricci scalar, and $|g|$ is the absolute value of the determinant of the euclidian metric tensor. The resulting theory is less singular than conventional quantum gravity. The action density of such a theory is a finite constant at zero curvature. So for $R = 0$, we can adjust θ and fit the density of dark energy at the classical level. One might be able to use

$$L_{sE}' = \frac{1}{8\pi G_N^2} \left(\frac{\cosh^2\theta}{\sqrt{1 - G_N \, R_E}} + \frac{\sinh^2\theta}{\sqrt{1 + G_N \, R_E}} \right) \sqrt{|g|} \tag{18}$$

to make the energy density of the ground state actually finite and equal to that of dark energy for an adjusted angle θ.

ACKNOWLEDGMENTS

I am grateful to Carl Bender, Ginette Cahill, Fred Cooper, Franco Giuliani, Roy Glauber, Gary Herling, and David Waxman for helpful conversations.

---

[1] S. Boettcher and C. Bender, J. Math. Phys. **31(11)**, 2579 (1990).

[2] K. Cahill, Phys. Rev. D (2013), (in press), arXiv:1212.0290 [hep-th].

[3] E. Komatsu, K. M. Smith, J. Dunkley, C. L. Bennett, B. Gold, G. Hinshaw, N. Jarosik, D. Larson, M. R. Nolta, L. Page, D. N. Spergel, M. Halpern, R. S. Hill, A. Kogut, M. Limon, S. S. Meyer, N. Odegard, G. S. Tucker, J. L. Weiland, E. Wollack, , and E. L. Wright, ApJS **192(2)** (2011), http://dx.doi.org/10.1088/0067-0049/192/2/18.

[4] K. Cahill, *Physical Mathematics* (Cambridge University Press, 2013) p. 643.

[5] K. G. Wilson, Phys. Rev. D **10**, 2445 (1974).

[6] M. Creutz, Phys. Rev. Lett. **43**, 553 (1979); **45**, 313 (1980).